Research Update Information

Clinical findings on Skin Health Regarding Phycocyanin, the Natural Blue Pigment from Spirulina, and its Possible Mechanisms of Action*

What is Linablue®, Phycocyanin?

EARTHRISE

Linablue[®] is a natural blue colorant, which contains phycocyanin and Is manufactured by Earthrise Nutritionals in Southern California, USA. Linablue[®] is produced through the extraction process from spirulina under the controlled facility with current Good Manufacturing Practice (cGMP), International Organization for Standardization (ISO 9001), and Food Safety System (FSSC 22000). Linablue[®] is non-GMO, Kosher and Halal certified. Phycocyanin is one of the main compounds in spirulina, and a variety of research regarding its chemistry and health benefits, e.g. immune, cardiovascular and antioxidant properties have been reported around the world. There are more than more than 2,500 publications according to the Public Medine (NIH database) search as of February, 2024.



Beta-carotene Chlorophyll Phycocyanin

CLINICAL STUDY SUMMARY

[Clinical study design] Randomized Double-blind Placebo-controlled
[Subjects] Healthy female: aged between 20 to 65 (n=96).
[Dosage] Phycocyanin 410 mg, including C-phycocyanin: 300 mg and allo-phycocyanin: 110 mg
[Duration] 8 wks
[Outcome] 1) Transepidermal water loss (TEWL) levels at 8 wks ↓ (P < 0.05).

2) VAS scores: skin elasticity and skin gloss at 4 and 8 wks and skin moisture at 8 wks ↑ (P < 0.05)





Reference

- 1) Imai, Y. et al. (2019) Phycocyanin reces transepidermal water loss and improves skin barrier function in healthy subjects. *Jpn Pharmacol. Ther.*, 47(11): 1833-40.
- 2) Kim, K.M. (2018) Phycocyanin Protects Against UVB-induced Apoptosis Through the PKC α/βII-Nrf-2/HO-1 Dependent Pathway in Human Primary Skin Cells. *Molecules* 23(2):478
- Jang, Y.A. and Kim, BA. (2021) Protective Effect of Spirulina-Derived C-Phycocyanin against Ultraviolet B-Induced Damage in HaCaT Cells. *Medicina (Kaunas)*. 57(3):273.
- 4) Zhou, Y. et al (2023). The anti-photoaging effect of C-phycocyanin on ultraviolet B-irradiated BALB/c-nu mouse skin. *Front Bioeng Biotechnol.* 11: 1229387.
- 5) Madhyastha, H. K. at al (2008) uPA dependent and independent mechanisms of wound healing by C-phycocyanin. *J. Cell. Mol. Med.* 12 (6B): 2691-2703.
- 6) Gur, C.S. et al., (2013) In vitro and in vivo investigation of the wound healing effect of crude spirulina extract and c-phycocyanin. *J. Med. Plants Res.* 7: 425-433.
- 7) Wu, L.C. et al., (2011) Antimelanogenic effect of c-phycocyanin through modulation of tyrosinase expression by upregulation of ERK and downregulation of p38 MAPK signaling pathways. *J. Biomed Sci.* 18:74

* These statements have not been evaluated by the Food and Drug Administration. These products are not intended to diagnose, treat, cure or prevent any disease.

Earthrise Nutritionals LLC

113 East Hoober Rd., Calipatria, CA 92233 Tel: (949) 623-0980, Email: information@earthrise.com Published in February, 2024